

# Subvarieties of general type on a general projective hypersurface

Gianluca Pacienza

**Abstract.** We study subvarieties of a general projective degree  $d$  hypersurface  $X_d \subset \mathbf{P}^n$ . Our main theorem, which improves previous results of L. Ein and C. Voisin, implies in particular the following sharp corollary: any subvariety of a general hypersurface  $X_d \subset \mathbf{P}^n$ , for  $n \geq 6$  and  $d \geq 2n - 2$ , is of general type.

## 1 Introduction

Let  $X_d \subset \mathbf{P}^n$  be a general (in the countable Zariski topology) complex projective hypersurface of degree  $d$ . The study of the geometry of  $k$ -dimensional subvarieties of  $X_d$  in terms of  $k$ ,  $n$  and  $d$  has received much attention in the last 15 years (see [C1], [E1] and [E2], [X1] and [X2], [V1] and [V2], [CL], [CLR], [P], [C2], [CR]). In particular this study is related to the hyperbolicity of the hypersurface  $X_d \subset \mathbf{P}^n$ . Recall that a compact complex manifold  $M$  is said to be *hyperbolic* (in the sense of Brody or Kobayashi) if there are no nonconstant entire holomorphic maps  $f : \mathbb{C} \rightarrow M$ . S. Lang conjectured (cf. [L], Conjecture 5.6) that, in the case of a projective variety  $V$ , the notion of hyperbolicity has an *algebraic* characterization, namely  $V$  is hyperbolic if and only if any subvariety  $Y$  of  $V$  is of general type (that is, if  $Y$  is smooth, some multiple of the canonical bundle of  $Y$  gives a projective embedding of a non-empty Zariski open subset of  $Y$ . If  $Y$  is singular, then it is said to be of general type if some desingularization of  $Y$  has this property). Notice that if any subvariety  $Y$  of  $V$  is of general type, then in particular  $V$  does not contain rational curves or abelian subvarieties - a condition which is of course implied by the hyperbolicity. In this paper we focus our attention

---

*2000 Mathematics Subject Classification.* Primary 14J70, 14K12, 14C99; Secondary 32Q45.

on the case of a general projective hypersurface  $X_d \subset \mathbf{P}^n$ , and give a sharp bound on its degree  $d$  in order to satisfy the algebraic property above that, conjecturally, should be equivalent to the hyperbolicity of  $X_d$ .

This problem has been studied by L. Ein in [E1] and [E2], where, generalizing a previous result by H. Clemens [C1], he proves in particular that whenever  $d \geq 2n - k$ , for  $n \geq 3$ , then any  $k$ -dimensional subvariety  $Y$  of the general  $X_d \subset \mathbf{P}^n$  has nonzero geometric genus, and if the inequality is strict then  $Y$  is of general type. Ein's result, which concerns more generally subvarieties of general complete intersections in an arbitrary smooth projective variety, has been improved by one, in the case of projective hypersurfaces, by C. Voisin ([V1], [V2]), who proves the following

**Theorem (Voisin).** *Let  $X_d \subset \mathbf{P}^n$  be a general hypersurface of degree  $d \geq 2n - k - 1$ , where  $k$  is an integer such that  $1 \leq k \leq n - 3$ . Then any  $k$ -dimensional subvariety  $Y$  of  $X$  has nonzero geometric genus, and if the inequality is strict then  $Y$  is of general type.*

Our main result is the following

**Theorem 1.1.** *Let  $X_d \subset \mathbf{P}^n$  be a general complex projective hypersurface of degree  $d$ . Let  $Y \subset X_d$  be a  $k$ -dimensional subvariety with a desingularization  $\tilde{Y} \xrightarrow{j} Y$  such that  $h^0(\tilde{Y}, K_{\tilde{Y}} \otimes j^* \mathcal{O}_{\mathbf{P}^n}(-1)) = 0$ . If the inequalities*

$$d - 1 \geq \max \left\{ \frac{7n - 3k - 3}{4}, \frac{3n - k + 1}{2} \right\} \quad (1)$$

and

$$\frac{d(d - 3)}{2} \geq 2n - k - 3 \quad (2)$$

are satisfied, then  $Y$  is contained in the locus covered by the lines of  $X$ .

Since the dimension of the locus spanned by the lines on a general hypersurface  $X_d$  is equal to  $(2n - 2) - d$ , it follows from Theorem 1.1 that for any subvariety  $Y$  of  $X_d \subset \mathbf{P}^n$ , whenever  $d \geq 2n - 2$  and  $n \geq 6$ , the canonical bundle of a desingularization  $j : \tilde{Y} \rightarrow Y$  is the sum of the effective divisor  $K_{\tilde{Y}} \otimes j^* \mathcal{O}_{\mathbf{P}^n}(-1)$  and of  $j^* \mathcal{O}_{\mathbf{P}^n}(1)$ , which is very ample on an open subset, so we obtain

**Corollary 1.2.** *Any subvariety of a general hypersurface  $X_d \subset \mathbf{P}^n$ , with  $d \geq 2n - 2$  and  $n \geq 6$ , is of general type.*

Corollary 1.2 is sharp, since general hypersurfaces of degree  $2n-3$  contain a finite number of lines (and, by [P], lines are the only rational curves allowed on the general  $X_{2n-3} \subset \mathbf{P}^n$ , for  $n \geq 6$ ).

The weird looking numerical hypotheses (1) and (2) of Theorem 1.1 are needed in order to control the positivity of the twisted exterior powers of the bundle  $M_{\mathbf{P}^n}^d$  (resp.  $M_G^d$ ) over  $\mathbf{P}^n$  (resp. over the Grassmannian of lines in  $\mathbf{P}^n$ ), that are defined in §2. This control will appear to be a crucial point along the proof (cf. Lemmas 3.1, 4.1 and 4.3). The proof of Theorem 1.1 makes use of the powerful variational approach introduced by C. Voisin in [V1] and [V2], and adopted by the author in [P] to study the geometry of subvarieties having geometric genus zero on a general hypersurface. These methods have been strengthened more recently by H. Clemens and Z. Ran (see [C2] and [CR]) to study in greater generality subvarieties  $Y$  of  $X$  with desingularizations  $j : \tilde{Y} \rightarrow Y$  verifying  $h^0(\tilde{Y}, K_{\tilde{Y}} \otimes j^* \mathcal{O}_{\mathbf{P}^n}(a)) = 0$ , for some integer  $a \geq 0$ .

The proof is naturally divided in two parts. First, following an idea that goes back to Voisin [V2], under a technical numerical hypothesis, it is possible to see that through each point of  $Y$  there is a line which intersects set-theoretically  $X$  in at most two points. Precisely we prove

**Proposition 1.3.** *Let  $X_d \subset \mathbf{P}^n$  be a general hypersurface, whose degree  $d$  verifies the numerical condition (1) of Theorem 1.1, and  $Y \subset X_d$  a subvariety of dimension  $k$  such that  $h^0(\tilde{Y}, K_{\tilde{Y}} \otimes j^* \mathcal{O}_{\mathbf{P}^n}(-1)) = 0$ , where  $j : \tilde{Y} \rightarrow Y$  is a desingularization. Then, for some  $r \geq 1$ ,  $Y$  is contained in the sublocus  $\Delta_{(r,d-r),X}$  of  $X_d$  defined as*

$$\Delta_{(r,d-r),X} := \{x \in X_d : \exists \text{ a line } \ell \text{ s.t. } \ell \cap X_d = r \cdot x + (d-r) \cdot x', x' \in X_d\}.$$

(This result, under different numerical hypothesis, can also be found in [CR]).

The second part of the proof of Theorem 1.1 deals with the study of the locus  $\Delta_{(r,d-r),X}$ . For this, I use two explicit desingularizations of  $\Delta_{(r,d-r),X}$ , whose canonical bundles are easily computable. Then, in both cases, the key point is the construction of a globally generated subbundle contained in the exterior powers of the (twisted) tangent bundle of the family of the desingularizations. This fact will allow us to obtain the following proposition which concludes the proof of Theorem 1.1:

**Proposition 1.4.** *Let  $X_d \subset \mathbf{P}^n$  be a general hypersurface of degree  $d$  verifying the numerical condition (2) of Theorem 1.1. Let  $Y \subset \Delta_{(r,d-r),X}$  be a subvariety of dimension  $k$ , and  $j : \tilde{Y} \rightarrow Y$  a desingularization such that  $h^0(\tilde{Y}, K_{\tilde{Y}} \otimes j^* \mathcal{O}_{\mathbf{P}^n}(-1)) = 0$ . Then  $Y$  is contained in the locus of lines of  $X_d$ .*

*Acknowledgements.* I would like to thank M. Zaidenberg and L. Manivel, who invited me, respectively on February 2001 and March 2002, at the Institut Fourier in Grenoble, where the preliminary version of this work was started and brought to end. My sincerest thanks to H. Clemens, who sent me the manuscript [C2] and kindly encouraged me when I communicated to him the idea of adapting some arguments contained in [C2] to strengthen my work. Finally, it is a pleasure to acknowledge the great influence of C. Voisin's articles [V1] and [V2] on my work.

## 2 Preliminaries

We will follow the notation already used in [P], which we recall below.

**Notation.**

$$S^d := H^0(\mathbf{P}^n, \mathcal{O}_{\mathbf{P}^n}(d));$$

$$S_x^d := H^0(\mathbf{P}^n, \mathcal{I}_x \otimes \mathcal{O}_{\mathbf{P}^n}(d));$$

$$N := h^0(\mathbf{P}^n, \mathcal{O}_{\mathbf{P}^n}(d)) = \dim S^d;$$

$\mathcal{X} \subset \mathbf{P}^n \times S^d$  will denote the universal hypersurface of degree  $d$ ;

$X_F \subset \mathbf{P}^n$  the fiber of the family  $\mathcal{X}$  over  $F \in S^d$ , i.e. the hypersurface defined by  $F$ .

Let  $U \rightarrow S^d$  be an étale map and  $\mathcal{Y} \subset \mathcal{X}_U$  a reduced and irreducible subscheme of relative dimension  $k$  (in the following, by abuse of notation, we will often omit the étale base change). Let  $\tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$  be a desingularization and  $\tilde{\mathcal{Y}} \xrightarrow{j} \mathcal{X}_U$  the natural induced map. We may obviously assume  $\mathcal{Y}$  invariant under some lift of the action of  $GL(n+1)$  (recall that  $g \in GL(n+1)$  acts on the product  $\mathbf{P}^n \times S^d$  as follows:  $g(x, F) = (g(x), (g^{-1})^*F)$ ). Let  $\pi : \mathcal{X} \rightarrow \mathbf{P}^n$  be the projection on the first component and  $T_{\mathcal{X}}^{vert}$  (resp.  $T_{\mathcal{Y}}^{vert}$ ) the vertical part of  $T_{\mathcal{X}}$  (resp.  $T_{\mathcal{Y}}$ ) w.r.t.  $\pi$ , i.e.  $T_{\mathcal{X}}^{vert}$  (resp.  $T_{\mathcal{Y}}^{vert}$ ) is the sheaf defined by

$$0 \rightarrow T_{\mathcal{X}}^{vert} \rightarrow T_{\mathcal{X}} \xrightarrow{\pi_*} T\mathbf{P}^n \rightarrow 0$$

(resp.  $0 \rightarrow T_{\mathcal{Y}}^{vert} \rightarrow T_{\mathcal{Y}} \xrightarrow{\pi_*} T\mathbf{P}^n$ ).

**Remark 2.0.1.** Let  $\mathcal{Y}$  be a subscheme of  $\mathcal{X} \subset \mathbf{P}^n \times S^d$  of relative dimension  $k$  and invariant under the action of  $GL(n+1)$ . Then:

(i) the map  $T\mathcal{Y} \xrightarrow{\pi_*} T\mathbf{P}^n$  is surjective and hence

$$\text{codim}_{T_{\mathcal{X},(y,F)}^{\text{vert}}} T_{\mathcal{Y},(y,F)}^{\text{vert}} = \text{codim}_{\mathcal{X}} \mathcal{Y} = n - k - 1.$$

(ii)  $T_{\mathcal{Y},(y,F)}^{\text{vert}}$  contains the vertical part of the tangent space to the orbit of the point  $(y, F)$  under the action of  $GL(n+1)$ , i.e.

$$T_{\mathcal{Y},(y,F)}^{\text{vert}} \supset \langle S_y^1 \cdot J_F^{d-1}, F \rangle,$$

where  $J_F^{d-1}$  is the Jacobian ideal of  $F$ .

Let  $d$  be a positive integer. Consider the bundle  $M_{\mathbf{P}^n}^d$  defined by the exact sequence

$$0 \rightarrow M_{\mathbf{P}^n}^d \rightarrow S^d \otimes \mathcal{O}_{\mathbf{P}^n} \xrightarrow{ev} \mathcal{O}_{\mathbf{P}^n}(d) \rightarrow 0, \quad (3)$$

whose fiber at a point  $x$  identifies by definition to  $S_x^d$ . From the inclusion  $\mathcal{X} \hookrightarrow \mathbf{P}^n \times S^d$  we get the exact sequence

$$0 \rightarrow T\mathcal{X}|_{X_F} \rightarrow T\mathbf{P}^n|_{X_F} \oplus (S^d \otimes \mathcal{O}_{X_F}) \rightarrow \mathcal{O}_{X_F}(d) \rightarrow 0,$$

which combined with (3) gives us

$$0 \rightarrow M_{\mathbf{P}^n|_{X_F}}^d \rightarrow T\mathcal{X}|_{X_F} \rightarrow T\mathbf{P}^n|_{X_F} \rightarrow 0. \quad (4)$$

In other words  $M_{\mathbf{P}^n|_{X_F}}^d$  identifies to the vertical part of  $T\mathcal{X} \otimes \mathcal{O}_{X_F}$  with respect to the projection to  $\mathbf{P}^n$ .

Let  $G := \text{Grass}(1, n)$  be the Grassmannian of lines in  $\mathbf{P}^n$ ,  $\mathcal{O}_G(1)$  the line bundle on  $G$  giving its Plücker polarization, and  $\mathcal{E}_d$  be the  $d^{\text{th}}$ -symmetric power of the dual of the tautological subbundle on  $G$ . Recall that the fibre of  $\mathcal{E}_d$  at a point  $[\ell]$  is, by definition, given by  $H^0(\ell, \mathcal{O}_\ell(d))$ .

Let  $M_G^d$  be the vector bundle on  $G$  defined as the kernel of the evaluation map:

$$0 \rightarrow M_G^d \rightarrow S^d \otimes \mathcal{O}_G \rightarrow \mathcal{E}_d \rightarrow 0.$$

Notice that the fiber of  $M_G^d$  at a point  $[\ell]$  is equal to  $H^0(\mathcal{I}_\ell(d))$ .

The bundles  $M_{\mathbf{P}^n}^d$  and  $M_G^d$  verify the following positivity properties, that will be often used in what follows:

**Lemma 2.1.** (i)  $M_{\mathbf{P}^n}^d \otimes \mathcal{O}_{\mathbf{P}^n}(1)$  is generated by its global sections;  
(ii)  $M_G^d \otimes \mathcal{O}_G(1)$  is generated by its global sections.

*Proof.* See, for instance, [P]. □

### 3 A first reduction

Let  $Y_F \subset X_F$  be a general ( $k$ -dimensional) fiber of the subfamily  $\mathcal{Y} \subset \mathcal{X}_U$ , and  $\tilde{Y}_F \xrightarrow{j} Y_F$  its desingularization. By abuse of notation, we will often write  $K_{\tilde{Y}_F}(-1)$  instead of  $K_{\tilde{Y}_F} \otimes j^* \mathcal{O}_{\mathbf{P}^n}(-1)$ .

Recall now the isomorphisms:

- (i)  $\Omega_{\tilde{Y}_F}^{N+k} \cong K_{\tilde{Y}_F}$ ;
- (ii)  $(\wedge^{n-1-k} T\mathcal{X}_U|_{X_F}) \otimes K_{X_F} \cong \Omega_{\mathcal{X}_U|_{X_F}}^{N+k}$ .

Set  $c := n - 1 - k = \text{codim}_{X_F} Y_F$ . Using (i) and (ii), from the natural morphism  $\Omega_{\mathcal{X}_U}^1 \rightarrow \Omega_{\tilde{Y}_F}^1$ , we get a map

$$(\wedge^c T\mathcal{X}_U|_{X_F}) \otimes K_{X_F} \cong \Omega_{\mathcal{X}_U|_{X_F}}^{N+k} \rightarrow \Omega_{\tilde{Y}_F}^{N+k} \cong K_{\tilde{Y}_F}, \quad (5)$$

and hence, after tensoring by  $\mathcal{O}_{X_F}(-1)$ ,

$$(\wedge^c T\mathcal{X}_U|_{X_F}) \otimes K_{X_F}(-1) \longrightarrow K_{\tilde{Y}_F}(-1). \quad (6)$$

Now taking global sections in (6) and using (4) we have the following commutative diagram

$$\begin{array}{ccc} H^0(\wedge^c T\mathcal{X}_U|_{X_F} \otimes K_{X_F}(-1)) & \longrightarrow & H^0(K_{\tilde{Y}_F}(-1)) \\ \cup & \nearrow & \\ H^0(\wedge^c M_{\mathbf{P}^n}^d|_{X_F} \otimes K_{X_F}(-1)) & & \end{array} \quad (7)$$

By hypothesis, we have that the composite map in (7) is identically zero. This, by the  $GL(n+1)$ -invariance of  $\mathcal{Y}$ , implies that  $T_{\mathcal{Y},(y,F)}^{vert}$  is then contained in the base locus of  $H^0(\wedge^c M_{\mathbf{P}^n}^d|_{X_F} \otimes K_{X_F}(-1))$ , considered as the space of sections of a line bundle on the Grassmannian of codimension  $c$  subspaces of  $T^{vert}\mathcal{X}|_{X_F}$ .

The generalization presented in [C2] of the variational approach introduced by C. Voisin in [V1] and [V2], and applied by the author in [P], starts with a sharp algebraic study of the base locus of the bundles  $\wedge^c M_{\mathbf{P}^n}^d(b)$ . Precisely, we will use the following

**Lemma 3.1. ([C2]).** *Suppose  $d$  verifies the numerical condition (1) of Theorem 1.1. Let  $A$  be a codimension  $c$  subspace of  $S_x^d = (T_{X,(x,F)})^{vert}$  which is in the base locus of  $H^0(\wedge^c M_{\mathbf{P}^n}^d(d-n-2))$ . Then there exists a line  $\ell_A$  passing through  $x$  such that*

$$rk \frac{A}{A \cap H^0(\mathcal{I}_{\ell_A}(d))} \leq n+1. \quad (8)$$

*Proof.* In [C2] there is the study of the base locus of  $H^0(\wedge^c M_{\mathbf{P}^n}^d(d-n-1+a))$ , for  $a \geq 0$  (this is the point where Lemma 2.1, (i) is used). Here we simply remark that the arguments presented in [C2] also apply to the  $a = -1$  case. For the reader's convenience we outline the idea of the proof (for the details see [C2]). The main point in [V2] and [P] was to produce, by Koszul complexes' techniques, *explicit* global sections of the bundle  $\wedge^c M_{\mathbf{P}^n}^d(b)$ , for the integers  $b, c, d$  considered there. This was used to deduce that, for a generic polynomial  $P \in S^{d-1}$ , the multiplication map

$$\begin{aligned} m_{P,A} : S_x^1 &\rightarrow S_x^d/A \\ L &\mapsto L \cdot P \text{ mod } A \end{aligned}$$

has rank one. H. Clemens considers more generally in [C2] the smallest integer  $s \geq 0$  such that rank of the multiplication map

$$\begin{aligned} m_{P,A,s} : S_x^1 &\rightarrow S_x^d/(A + Q_1 \cdot S_x^1 + \dots + Q_s \cdot S_x^1) \\ L &\mapsto L \cdot P \text{ mod } (A + Q_1 \cdot S_x^1 + \dots + Q_s \cdot S_x^1) \\ P, Q_1, \dots, Q_s &\text{ are generic polynomials in } S^{d-1} \end{aligned}$$

is one. Then, as in [V2] and [P], an infinitesimal argument applies. Namely, recall that if  $V$  and  $W$  are vector spaces, and  $Z_k := \{\phi \in Hom(V, W) : rank \phi \leq k\}$ , then

$$T_{Z_k, \phi} = \{\psi \in Hom(V, W) : \psi(ker \phi) \subset Im \phi\}. \quad (9)$$

Applying this to the map  $m_{P,A,s}$ , we obtain that, for any  $R \in S^{d-1}$ ,

$$R \cdot Ker m_{P,A,s} \text{ mod } (A + Q_1 \cdot S_x^1 + \dots + Q_s \cdot S_x^1) \subset Im m_{P,A,s}$$

i.e.

$$H^0(\mathcal{I}_{\ell_A}(d)) \subset A + Q_1 \cdot S_x^1 + \dots + Q_s \cdot S_x^1 + P \cdot S_x^1,$$

where  $\ell_A$  is the line determined by  $Ker m_{P,A,s}$ . To complete the proof it remains to verify that the line is independent of the choice of the polynomials, and that, under the hypothesis (1), the integer  $s$  is such that  $A$  and  $H^0(\mathcal{I}_{\ell_A}(d))$  satisfy (8).  $\square$

Since the map (7) vanishes, Lemma 3.1 applies to the tangent space  $T_{\mathcal{Y},(y,F)}^{vert}$ . Then, at a generic point  $(y, F) \in \mathcal{Y}$ , the tangent space  $T_{\mathcal{Y},(y,F)}^{vert}$  contains a subspace  $T \subset H^0(\mathcal{I}_{\ell(y,F)}(d))$ , where  $\ell(y,F)$  is a line through  $y$  and  $T$  satisfies

$$rk \frac{T_{\mathcal{Y},(y,F)}^{vert}}{T} \leq n + 1. \quad (10)$$

We now verify an easy fact:

**Lemma 3.2.** *The tangent space  $T_{\mathcal{Y},(y,F)}^{vert}$  cannot contain two subspaces  $T$  and  $T'$  of ideals of different lines  $\ell \neq \ell'$  and verifying (10).*

*Proof.* Indeed, if this were the case, by the surjectivity of

$$H^0(\mathcal{I}_\ell(d)) \oplus H^0(\mathcal{I}_{\ell'}(d)) \twoheadrightarrow S_y^d,$$

and the numerical condition (10), then  $T_{\mathcal{Y},(y,F)}^{vert}$  would contain a subspace of  $S_y^d$  of codimension at most  $2(n + c + 1 - d)$ . Now, by remark 2.0.1, (i), we have

$$c = \text{codim}_{T_{\mathcal{X},(y,F)}^{vert}} T_{\mathcal{Y},(y,F)}^{vert} = \text{codim}_{\mathcal{X}} \mathcal{Y} \leq 2(n + c + 1 - d)$$

which is equivalent to

$$d \leq \frac{3n - k}{2},$$

and the last inequality is impossible because of (1).  $\square$

Then, we can consider the distribution  $\mathcal{T} \subset T_{\mathcal{Y}}^{vert}$ , pointwise given by the  $T$ 's. This distribution turns out to have the following properties

**Proposition 3.3.** *The distribution  $\mathcal{T} \subset T_{\mathcal{Y}}^{vert}$  is integrable and the natural map  $\phi : \mathcal{Y} \rightarrow G(1, n)$ , associating to  $(y, F)$  the line determined by  $T \subset H^0(\mathcal{I}_{\ell(y,F)}(d))$ , is constant along the leaves of the corresponding foliation.*

*Proof.* The proof goes along the lines of [V2], lemma 3 and 4, and of [P], lemma 3.3. For the detailed proof in the general case, see [C2]. Again, for the reader's convenience, we sketch it below. Consider the bracket map

$$\Psi : \bigwedge^2 \mathcal{T} \rightarrow T_{\mathcal{Y}}^{vert} / \mathcal{T} \subset T_{\mathcal{X}}^{vert} |_{\mathcal{Y}} / \mathcal{T},$$

which is given at the point  $(y, F)$  by

$$\psi : \wedge^2 T_{\ell(y,F)} \rightarrow T_{\mathcal{Y},(y,F)}^{vert} / T_{\ell(y,F)} \hookrightarrow H^0(\mathcal{O}_{\ell(y,F)}(d)(-y)).$$

Now, choose coordinates on  $\mathbf{P}^n$  such that  $\ell := \ell_{(y,F)} = \{X_2 = \dots = X_n = 0\}$  and  $y = [1, 0, \dots, 0]$ . Note that, since  $y \in \ell$ ,  $\phi_*(T_{\mathcal{Y},(y,F)}^{vert})$  is contained in  $H^0(N_{\ell/\mathbf{P}^n}(-y))$  ( $\phi_*$  is the differential of  $\phi$  at  $(y, F)$ ). One verifies that

$$\psi(A \wedge B) = A \cdot \phi_*(B) - B \cdot \phi_*(A), \quad A, B \in T_{\ell_{(y,F)}}, \quad (11)$$

where the bilinear map  $(a, b) \mapsto a \cdot b$  is explicitly given by

$$P \cdot (X_1 \sum_{i=2}^n b_i \frac{\partial}{\partial X_i}) = \sum_{i=2}^n b_i X_1 (\frac{\partial P}{\partial X_i})|_{\ell} \in H^0(\mathcal{O}_{\ell}(d)(-y)).$$

A key linear algebra lemma allows to prove that  $\phi_*$  is zero, so the proposition follows from (11) and from the Frobenius theorem.  $\square$

Using this we will prove, via the  $GL(n+1)$ -invariance of  $\mathcal{Y}$ , that  $F|_{\ell_{(y,F)}}$  has, set-theoretically, at most two zeroes.

*Proof of Proposition 1.3.* Let  $(y, F)$  be a general point of  $\mathcal{Y}$ . Let  $\ell$  be the line through  $y$  and  $T$  the subspace of  $H^0(\mathcal{I}_{\ell}(d))$  contained in  $T_{\mathcal{Y},(y,F)}^{vert}$  and verifying (10). Consider the following diagram

$$\begin{array}{ccccc} & & 0 & & 0 & (12) \\ & & \downarrow & & \downarrow & \\ 0 & \longrightarrow & T & \longrightarrow & H^0(\mathcal{I}_{\ell}(d)) & \\ & & \downarrow & & \downarrow & \\ 0 & \longrightarrow & T_{\mathcal{Y},(y,F)}^{vert} & \longrightarrow & T_{\mathcal{X},(y,F)}^{vert} = S_y^d & \\ & & \downarrow & & \downarrow & \\ & & H^0(\mathcal{O}_{\ell}(d)(-y)) & \equiv & H^0(\mathcal{O}_{\ell}(d)(-y)) & \\ & & & & \downarrow & \\ & & & & 0 & \end{array}$$

By (10) we have

$$\dim \operatorname{Im} (T_{\mathcal{Y},(y,F)}^{vert} \rightarrow H^0(\mathcal{O}_{\ell}(d)(-y))) \leq n+1 \quad (13)$$

On the other hand, by remark 2.0.1, (ii), the vertical tangent space  $T_{\mathcal{Y},(y,F)}^{vert}$  contains  $S_y^1 \cdot J_F^{d-1}$  and  $F$  itself. Take coordinates  $X_0, \dots, X_n$  on  $\mathbf{P}^n$  such that  $y = [1, 0, \dots, 0]$ , and  $\ell := \ell_{(y,F)} = \{X_2 = \dots = X_n = 0\}$ . Since  $\phi$  is constant along the leaves of the foliation, we can generically

choose a polynomial  $G$  in the leaf integrating  $T$  so that the  $(n-1)$ -elements  $X_1 \frac{\partial G}{\partial X_i}$ ,  $i \geq 2$ , are independent modulo the subspace

$$K := \langle G|_\ell, X_1 \left( \frac{\partial G}{\partial X_0} \right)|_\ell, X_1 \left( \frac{\partial G}{\partial X_1} \right)|_\ell \rangle \subset H^0(\ell, \mathcal{O}_\ell(d) \otimes \mathcal{I}_y),$$

which is uniquely determined by  $F|_\ell$  and hence is constant along the leaf integrating  $T$ . By (13), this implies  $\dim K \leq 2$ , that is

$$F|_\ell = \alpha X_1^r L^{d-r},$$

for some  $r \geq 1$ , and some linear form  $L$  on  $\ell$ . □

We are then led to study the locus  $\Delta_{(r,d-r),F}$ . This will be done in the last section.

## 4 The bicontact locus $\Delta_{(r,d-r),F}$

Let  $X_F \subset \mathbf{P}^n$  be a general hypersurface of degree  $d$  verifying (1), and  $Y_F \subset X_F$  a  $k$ -dimensional subvariety whose desingularization  $\tilde{Y}$  is such that  $h^0(\tilde{Y}, K_{\tilde{Y}}(-1)) = 0$ . Then, by Proposition 1.3, we know that  $Y_F$  is contained in  $\Delta_{(r,d-r),X_F} \subset X_F$ , the  $(2n-d)$ -dimensional subvariety of points  $x$  of  $X_F$  through which there is an osculating line  $\ell$  intersecting  $X_F$  at most at another point, *i.e.*  $\ell \cap X_F = r \cdot x + (d-r) \cdot x'$ ,  $x' \in X_F$ . In what follows we will write  $\Delta_{(r,d-r),F}$  instead of  $\Delta_{(r,d-r),X_F}$ . To prove our theorem, we study two explicit desingularizations of  $\Delta_{(r,d-r),F}$ , which have been used in [V2], that are both given in terms of the zero locus of a section of a vector bundle. Thus we compute, by adjunction, the canonical bundle of such a desingularization. Then, again, we adopt a variational approach and construct, in both cases, a subbundle contained in the exterior powers of the (twisted) tangent bundle to the family of the desingularizations. A positivity result, namely the global generation of this subbundle, allows us to conclude the proof.

**Case 1:**  $r \geq 2$  and  $d-r \geq 2$ .

Let  $G := Gr(1, n)$  be the Grassmannian of lines in  $\mathbf{P}^n$ . Let  $\mathcal{O}_G(1)$  be the line bundle on  $G$  which gives the Plücker embedding. Let  $Z$  be the blow-up along the diagonal  $\Delta$  of the product  $\mathbf{P}^n \times \mathbf{P}^n$  with projections:

$$\begin{array}{ccc} Z := Bl_\Delta \mathbf{P}^n \times \mathbf{P}^n & \xrightarrow{b} & \mathbf{P}^n \times \mathbf{P}^n \xrightarrow{p_2} \mathbf{P}^n \\ & & \downarrow p_1 \\ & & \mathbf{P}^n \end{array} \quad (14)$$

Consider the map

$$\begin{aligned} f : Z &\rightarrow Gr(1, n), \\ z &\mapsto \ell_z \end{aligned} \tag{15}$$

where  $\ell_z$  is the line determined by  $z$ . Let  $\tilde{p}_i := p_i \circ b$ , for  $i = 1, 2$ , and consider the line bundles on  $Z$  defined as follows:  $H_i := \tilde{p}_i^* \mathcal{O}_{\mathbf{P}^n}(1)$  and  $L := f^* \mathcal{O}_G(1)$ . The variety  $Z$  comes together with a projective bundle:  $\mathbf{P} \xrightarrow{\pi} Z$ , and we define  $\mathcal{E}_d := \pi_* \mathcal{O}_{\mathbf{P}}(d)$ . Notice that the fibre of  $\mathcal{E}_d$  at  $z$  is equal to  $H^0(\ell_z, \mathcal{O}_{\ell_z}(d))$ . Consider the line bundle  $\mathcal{L}_{r,d-r} \subset \mathcal{E}_d$ , whose fibre at  $z \in Z$  is given by the one dimensional space of polynomials  $P \in H^0(\ell_z, \mathcal{O}_{\ell_z}(d))$  vanishing at  $x$  to the order  $r$  and at  $y$  to the order  $d - r$ , where  $(x, y) = b(z) \in \mathbf{P}^n \times \mathbf{P}^n$ . Define  $\mathcal{F}_{r,d-r} := \mathcal{E}_d / \mathcal{L}_{r,d-r}$ . To any polynomial  $F \in S^d$  we can associate a section  $\sigma_F \in H^0(Z, \mathcal{E}_d)$ , whose value at a point  $z$  is exactly the polynomial  $F|_{\ell_z} \in \mathcal{E}_d|_z$ , and we will denote by  $\bar{\sigma}_F$  its image in  $H^0(Z, \mathcal{F}_{r,d-r})$ . Then we define  $\tilde{\Delta}_{(r,d-r),F} := V(\bar{\sigma}_F)$ . By construction, we have  $\tilde{p}_1(\tilde{\Delta}_{(r,d-r),F}) = \Delta_{(r,d-r),F}$ . Since  $\mathcal{F}_{r,d-r}$  is generated by its global sections, the variety  $\tilde{\Delta}_{(r,d-r),F}$  is smooth and of the right dimension. Moreover, since in the degree considered through a generic point of  $\Delta_{(r,d-r),F}$  there is just one  $r$ -osculating line, the map  $\tilde{p}_1$  is a desingularization of  $\Delta_{(r,d-r),F}$ . We will now recall how to compute the canonical bundle of  $\tilde{\Delta}_{(r,d-r),F}$ . As remarked in [V2], the Picard group of  $Z$  is generated by  $H_1$ ,  $H_2$  and  $L$ , the canonical class of  $Z$  is  $K_Z = -2H_1 - 2H_2 + (-n+1)L$ , and the class of  $\mathcal{L}_{(r,d-r),F}$  is given by  $rH_1 + (d-r)H_2$ . Therefore, by adjunction, we have

$$K_{\tilde{\Delta}_{(r,d-r),F}} = K_Z + c_1(\mathcal{F}_{r,d-r}) = (r-2)H_1 + (d-r-2)H_2 + \left(\frac{d(d-1)}{2} - n+1\right)L.$$

Consider now the bundles  $\mathcal{N}_Z^{r,d-r}$  and  $\mathcal{M}_Z^d$  on  $Z$  respectively defined by the two following exact sequences :

$$0 \rightarrow \mathcal{N}_Z^{r,d-r} \rightarrow S^d \otimes \mathcal{O}_Z \rightarrow \mathcal{F}_{r,d-r} \rightarrow 0, \tag{16}$$

$$0 \rightarrow \mathcal{M}_Z^d \rightarrow S^d \otimes \mathcal{O}_Z \rightarrow \mathcal{E}_d \rightarrow 0. \tag{17}$$

By definition we have

$$0 \rightarrow \mathcal{M}_Z^d \rightarrow \mathcal{N}_Z^{r,d-r} \rightarrow \mathcal{L}_{r,d-r} \rightarrow 0. \tag{18}$$

The needed positivity result is the following

**Lemma 4.1.** *If  $r \geq 3$ ,  $d - r \geq 2$  and*

$$\frac{d(d-1)}{2} - n \geq c - 1, \quad (19)$$

*then the bundle  $\wedge^c \mathcal{M}_{Z|\tilde{\Delta}_{(r,d-r),F}}^d \otimes K_{\tilde{\Delta}_{(r,d-r),F}}(-H_1)$  is generated by its global sections.*

*Proof.* Remark that  $\mathcal{M}_Z^d = f^* M_G^d$ . Hence, by lemma 2.1, (ii), the bundle

$$\begin{aligned} & \wedge^c \mathcal{M}_Z^d \otimes \det \mathcal{F}_{r,d-r} \otimes K_Z(-H_1) = \\ & = f^*(\wedge^c M_G^d) \otimes \mathcal{O}_Z((r-3)H_1 + (d-r-2)H_2 + (\frac{d(d-1)}{2} - n + 1)L) \\ & = f^*(\wedge^c M_G^d(c)) \otimes \mathcal{O}_Z((r-3)H_1 + (d-r-2)H_2 + (\frac{d(d-1)}{2} - n - c + 1)L), \end{aligned}$$

is globally generated under our numerical hypothesis, and the same holds for its restriction to  $\tilde{\Delta}_{(r,d-r),F}$ .  $\square$

Let now  $\Delta_{r,d-r} \subset \mathbf{P}^n \times S^d$  be the family of the  $\Delta_{(r,d-r),F}$ 's, and  $\tilde{\Delta}_{r,d-r} \subset Z \times S^d$  the family of the desingularizations. Let  $\mathcal{Y} \subset \tilde{\Delta}_{r,d-r}$  be a subscheme of relative dimension  $k$ , invariant under the action of  $GL(n+1)$ , and  $\tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$  a desingularization. Assume  $h^0(\tilde{\mathcal{Y}}_F, K_{\tilde{\mathcal{Y}}_F}(-H_1)) = 0$  and set  $c = n - 1 - k$ . Recall the isomorphisms

$$\wedge^c T\tilde{\Delta}_{r,d-r}|_{\tilde{\Delta}_{(r,d-r),F}} \otimes K_{\tilde{\Delta}_{(r,d-r),F}} \cong \Omega_{\tilde{\Delta}_{r,d-r}|\tilde{\Delta}_{(r,d-r),F}}^{N+k} \quad (20)$$

$$\Omega_{\tilde{\mathcal{Y}}|\tilde{\mathcal{Y}}_F}^{N+k} \cong K_{\tilde{\mathcal{Y}}_F} \quad (21)$$

and consider the natural map

$$\wedge^c T\tilde{\Delta}_{r,d-r}|_{\tilde{\Delta}_{(r,d-r),F}} \otimes K_{\tilde{\Delta}_{(r,d-r),F}} \cong \Omega_{\tilde{\Delta}_{r,d-r}|\tilde{\Delta}_{(r,d-r),F}}^{N+k} \rightarrow \Omega_{\tilde{\mathcal{Y}}|\tilde{\mathcal{Y}}_F}^{N+k} \cong K_{\tilde{\mathcal{Y}}_F}. \quad (22)$$

If we twist (22) by  $-H_1$ , then, by assumption, the induced map in cohomology

$$H^0(\wedge^c T\tilde{\Delta}_{r,d-r}|_{\tilde{\Delta}_{(r,d-r),F}} \otimes K_{\tilde{\Delta}_{(r,d-r),F}}(-H_1)) \rightarrow H^0(K_{\tilde{\mathcal{Y}}_F}(-H_1)) \quad (23)$$

is zero. Let  $T_{\tilde{\Delta}_{r,d-r}}^{vert}$  be the sheaf defined by

$$0 \rightarrow T_{\tilde{\Delta}_{r,d-r}}^{vert} \rightarrow T\tilde{\Delta}_{r,d-r} \rightarrow TZ \rightarrow 0.$$

Its restriction to  $\tilde{\Delta}_{(r,d-r),F}$  coincides with  $\mathcal{N}_Z^{r,d-r}|_{\tilde{\Delta}_{(r,d-r),F}}$ . Therefore, by (18) and lemma 4.1, we have constructed a sub-bundle

$$\wedge^c \mathcal{M}_Z^d|_{\tilde{\Delta}_{(r,d-r),F}} \otimes K_{\tilde{\Delta}_{(r,d-r),F}}(-H_1) \hookrightarrow \wedge^c T_{\tilde{\Delta}_{r,d-r}}^{vert}|_{\tilde{\Delta}_{(r,d-r),F}} \otimes K_{\tilde{\Delta}_{(r,d-r),F}}(-H_1),$$

which is generated by its global sections, under the numerical hypothesis of the lemma 4.1.

We conclude the case 1 with the following

**Proposition 4.2.** *Let  $F$  be a general polynomial of degree  $d$  verifying (19), with  $c = n - 1 - k$ . Suppose  $r \geq 2$  and  $d - r \geq 2$ . Let  $Y_F \subset \tilde{\Delta}_{(r,d-r),F}$  be a subvariety of dimension  $k$ , and  $j : \tilde{Y}_F \rightarrow Y_F$  a desingularization such that  $h^0(\tilde{Y}_F, K_{\tilde{Y}_F}(-j^*H_1)) = 0$ . Then  $Y_F$  is contained in the locus of lines of  $X_F$ .*

*Proof.* When no confusion is possible, we will omit in what follows the index  $(r, d - r)$  and simply set  $\Delta = \Delta_{r,d-r}$ . Suppose first  $r \geq 3$ . Let  $W \subset T_{\tilde{\Delta},(z,F)}$  be a codimension  $c$  subspace contained in the base locus of  $H^0(\wedge^c T_{\tilde{\Delta}|_{\tilde{\Delta}_F}} \otimes K_{\tilde{\Delta}_F}(-H_1))$ , considered as the space of sections of a line bundle on the Grassmannian of codimension  $c$  subspaces of  $T_{\tilde{\Delta}|_{\tilde{\Delta}_F}}$ . Then we must have

$$W^{vert} := W \cap \mathcal{N}_Z^{r,d-r}|_z \subset \mathcal{M}_Z^d|_z. \quad (24)$$

Indeed, if this were not the case, we would have  $\text{codim}_{\mathcal{M}_Z^d|_z} \bar{W} = c$ , where  $\bar{W} := W \cap \mathcal{M}_Z^d|_z$ . Then consider the following commutative diagram:

$$\begin{array}{ccc} H^0(\wedge^c \mathcal{M}_{Z|\tilde{\Delta}_F}^d \otimes K_{\tilde{\Delta}_F}(-H_1)) & \xrightarrow{\quad} & H^0(\wedge^c T_{\tilde{\Delta}|_{\tilde{\Delta}_F}} \otimes K_{\tilde{\Delta}_F}(-H_1)) \\ \downarrow ev & & \downarrow ev \\ (\wedge^c \mathcal{M}_{Z|\tilde{\Delta}_F}^d \otimes K_{\tilde{\Delta}_F}(-H_1))|_z & \xrightarrow{\quad} & (\wedge^c T_{\tilde{\Delta}|_{\tilde{\Delta}_F}} \otimes K_{\tilde{\Delta}_F}(-H_1))|_z \\ & \searrow \langle \cdot, \bar{W} \rangle & \downarrow \langle \cdot, W \rangle \\ & & \mathbb{C} \end{array} \quad (25)$$

( $ev$  is the evaluation of the sections at the point  $z$ , and  $\langle \cdot, W \rangle$  is the contraction defined by the subspace  $W$ ). Since  $W$  belongs to the base locus of  $H^0(\wedge^c T_{\tilde{\Delta}|_{\tilde{\Delta}_F}} \otimes K_{\tilde{\Delta}_F}(-H_1))$ , then the composite map  $\langle \cdot, W \rangle \circ ev$  is zero, and so would be  $\langle \cdot, \bar{W} \rangle \circ ev$ . But this is absurd, because, by Lemma 4.1, the bundle  $\wedge^c \mathcal{M}_{Z|\tilde{\Delta}_F}^d \otimes K_{\tilde{\Delta}_F}(-H_1)$  is generated by its global sections.

Let then  $\mathcal{Y} \subset \tilde{\Delta}$  be a subvariety, which is stable under the action of  $GL(n+1)$  and of relative codimension  $c$ . Assume moreover that the restriction map (23) is zero. By (24),  $T_{\mathcal{Y},(z,F)}^{vert}$  is contained in

$$\mathcal{M}_{Z|z}^d = \{G \in S^d : G|_{\ell_z} = 0\}. \quad (26)$$

On the other hand, by Remark 2.0.1, (ii),  $T_{\mathcal{Y},(z,F)}^{vert}$  contains  $F$  itself. So by (26) we have that  $F|_{\ell_z} = 0$  for every point  $z \in Y_F$ , i.e.  $Y_F$  is contained in the subvariety covered by the lines contained in  $X_F$ .

If  $r = 2$ , we can consider the natural isomorphism

$$\tilde{\Delta}_{(r,d-r),F} \xrightarrow{\sim} \tilde{\Delta}_{(d-r,r),F} \quad (27)$$

sending a point  $z \in \tilde{\Delta}_{(r,d-r),F}$ , with  $b(z) = (x, x')$ , to a point  $w \in \tilde{\Delta}_{(d-r,r),F}$  with  $b(w) = (x', x)$ , where  $x$  and  $x'$  are points on  $X_F$  linked by the condition

$$\exists \text{ a line } \ell \text{ s.t. } \ell \cap X_F = r \cdot x + (d-r) \cdot x'.$$

Since  $r = 2$  implies  $d - r \geq 3$ , from what we have done before it follows that the conclusion is true for  $\tilde{\Delta}_{(d-r,r),F}$ , and so, by (27), the same holds for  $\tilde{\Delta}_{(r,d-r),F}$ .  $\square$

**Case 2:**  $r = 1$  or  $d - r = 1$ .

Suppose for instance  $d - r = 1$ . Let  $\Gamma \subset \mathbf{P}^n \times Gr(1, n)$  be the incidence variety, and  $p$  and  $q$  the projections on the two factors. Let  $\pi : \mathbf{P} \rightarrow \Gamma$  be the pull-back of the universal  $\mathbf{P}^1$ -bundle over  $Gr(1, n)$  and  $\tau$  the natural section of  $\pi$ . Consider the bundle  $\mathcal{E}_d := \pi_* \mathcal{O}_{\mathbf{P}}(d)$  over  $\Gamma$ , and its rank 2 subbundle  $\mathcal{K} \subset \mathcal{E}_d$  such that its fiber at a point  $(x, \ell)$  is given by the polynomials  $P \in H^0(\mathcal{O}_{\ell}(d))$  vanishing to the order at least  $(d - 1)$  at  $x$ . Consider the line bundle  $\mathcal{L}_1$  defined by

$$0 \rightarrow \mathcal{L}_1 \rightarrow \mathcal{E}_1 \rightarrow \tau^* \mathcal{O}_{\mathbf{P}}(1) =: H \rightarrow 0.$$

Then  $\mathcal{K} \cong \mathcal{L}_1^{d-1} \otimes \mathcal{E}_1$ . Let  $\mathcal{F}_d$  be the quotient  $\mathcal{E}_d/\mathcal{K}$ . As in Case 1, to any  $F \in S^d$  we can associate a global section  $\sigma_F$  of  $\mathcal{F}_d$ . By definition  $p(V(\sigma_F)) = \Delta_{(d-1,1),F}$ . As before, since the bundle  $\mathcal{F}_d$  is generated by the sections  $\sigma_F$ , we have that  $V(\sigma_F)$  is smooth of the right dimension, for a general  $F$ , and it is easy to verify that  $p : V(\sigma_F) \rightarrow \Delta_{(d-1,1),F}$  is a desingularization. Then we define  $\tilde{\Delta}_{(d-1,1),F} := V(\sigma_F)$ . Hence, using the adjunction formula, we can compute the canonical bundle of  $\tilde{\Delta}_{(d-1,1),F}$  as the restriction to  $\tilde{\Delta}_{(d-1,1),F}$  of the following line bundle:

$$K_{Gr(1,n)} + c_1(\mathcal{F}_d) = (2d - 4)H + \left(\frac{d(d+1)}{2} - n - 1 - 2(d-1)\right)L. \quad (28)$$

Consider the bundles  $\mathcal{N}_{\Gamma}^d$  and  $\mathcal{M}_{\Gamma}^d$  on  $\Gamma$  respectively defined by the two following exact sequences :

$$0 \rightarrow \mathcal{N}_{\Gamma}^d \rightarrow S^d \otimes \mathcal{O}_{\Gamma} \rightarrow \mathcal{F}_d \rightarrow 0, \quad (29)$$

$$0 \rightarrow \mathcal{M}_\Gamma^d \rightarrow S^d \otimes \mathcal{O}_\Gamma \rightarrow \mathcal{E}_d \rightarrow 0. \quad (30)$$

From the definitions it follows that we have

$$0 \rightarrow \mathcal{M}_\Gamma^d \rightarrow \mathcal{N}_\Gamma^d \rightarrow \mathcal{K} \rightarrow 0. \quad (31)$$

The positivity result we will need this time is the following

**Lemma 4.3.** *If*

$$\frac{d(d+1)}{2} - n - 1 - 2(d-1) \geq c - 1, \quad (32)$$

*then the bundle  $\wedge^c \mathcal{M}_{\Gamma|\tilde{\Delta}_{(d-1,1),F}}^d \otimes K_{\tilde{\Delta}_{(d-1,1),F}}(-H)$  is generated by its global sections.*

*Proof.* Use (28) and lemma 2.1, (ii). □

Let now  $\Delta_{d-1,1} \subset \mathbf{P}^n \times S^d$  be the family of the  $\Delta_{(d-1,1),F}$ 's, and  $\tilde{\Delta}_{d-1,1} \subset \Gamma \times S^d$  the family of the desingularizations. Let  $\mathcal{Y} \subset \tilde{\Delta}_{r,d-r}$  be a subscheme of relative dimension  $k$ , invariant under the action of  $GL(n+1)$ , and  $\tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$  a desingularization. Consider the sheaf  $T_{\tilde{\Delta}_{d-1,1}}^{vert}$  defined by

$$0 \rightarrow T_{\tilde{\Delta}_{d-1,1}}^{vert} \rightarrow T_{\tilde{\Delta}_{r,d-r}} \rightarrow T_\Gamma \rightarrow 0,$$

and remark that its restriction to  $\tilde{\Delta}_{(d-1,1),F}$  coincides with  $\mathcal{N}_{\Gamma|\tilde{\Delta}_{(d-1,1),F}}^d$ . If we assume  $h^0(\tilde{Y}_F, K_{\tilde{Y}_F}(-H)) = 0$  and set  $c = n - 1 - k$ , we have that the natural adjunction map,

$$H^0(\wedge^c \mathcal{M}_{\Gamma|\tilde{\Delta}_{(d-1,1),F}}^d \otimes K_{\tilde{\Delta}_{(d-1,1),F}}(-H)) \rightarrow H^0(\tilde{Y}_F, K_{\tilde{Y}_F}(-H)), \quad (33)$$

that we can construct thanks to (31), is obviously zero.

The last step will be the proof of the following

**Proposition 4.4.** *Let  $F$  be a general polynomial of degree  $d$  verifying (32). Let  $Y_F \subset \tilde{\Delta}_{(d-1,1),F}$  be a subvariety of dimension  $k$ , and  $j : \tilde{Y}_F \rightarrow Y_F$  a desingularization such that (33) vanishes. Then  $Y_F$  is contained in the locus of lines of  $X_F$ .*

*Proof.* Recall that  $F \in T^{vert}\mathcal{Y}|_{(x,\ell,F)}$ . We claim that

$$F \in \mathcal{M}_{\Gamma|_{(x,\ell)}}^d.$$

Indeed if  $F \in \mathcal{K}|_{(x,\ell)}$ , then we have the surjection

$$T^{vert}\mathcal{Y}|_{(x,\ell,F)} \twoheadrightarrow \mathcal{K}|_{(x,\ell)}. \quad (34)$$

(This follows from the fact that if  $F \in \mathcal{K}|_{(x,\ell)}$  then

$$\langle S_x^1 \cdot J_F^{d-1}, F \rangle \twoheadrightarrow \mathcal{K}|_{(x,\ell)},$$

plus Remark 2.0.1, (ii)). Then, by (34), we have that

$$\text{codim}_{\mathcal{M}_{\Gamma|_{(x,\ell)}}^d} T^{vert}\mathcal{Y}|_{(x,\ell,F)} = \text{codim}_{X_F} Y_F = c.$$

As in Proposition 4.2, we can now use the commutative diagram

$$\begin{array}{ccc} H^0(\wedge^c \mathcal{M}_{\Gamma|\tilde{\Delta}_F}^d \otimes K_{\tilde{\Delta}_F}(-H)) & \hookrightarrow & H^0(\wedge^c T\tilde{\Delta}|_{\tilde{\Delta}_F} \otimes K_{\tilde{\Delta}_F}(-H)) \\ \downarrow ev & & \downarrow ev \\ (\wedge^c \mathcal{M}_{\Gamma|\tilde{\Delta}_F}^d \otimes K_{\tilde{\Delta}_F}(-H))|_{(x,\ell)} & \hookrightarrow & (\wedge^c T\tilde{\Delta}|_{\tilde{\Delta}_F} \otimes K_{\tilde{\Delta}_F}(-H))|_{(x,\ell)} \\ & \searrow \langle \cdot, \tilde{W} \rangle & \downarrow \langle \cdot, W \rangle \\ & & \mathbb{C} \end{array} \quad (35)$$

and deduce from it, together with Lemma 4.3 and Remark 2.0.1, (ii), that what we claim holds, *i.e.*

$$F|_{\ell} = 0.$$

□

The numerical condition (32), which, for  $c = n - 1 - k$ , becomes

$$\frac{d(d-3)}{2} \geq 2n - k - 3,$$

implies (19). Thus, combining the two propositions 4.2 and 4.4, the proof of our main theorem is completed. □

## References

- [C1] H. Clemens, *Curves in generic hypersurfaces*, Ann. Sci. École Norm. Sup. **19** (1986), 629–636.

- [C2] H. Clemens, *Limiting the genus of subvarieties of generic hypersurfaces*, Comm. in Algebra, Special issue in honour of S. Kleiman, to appear.
- [CL] L. Chiantini, A. F. Lopez, *Focal loci of families and the genus of curves on surfaces*, Proc. Amer. Math. Soc. **127** (1999), no. 12, 3451–3459.
- [CLR] L. Chiantini, A. F. Lopez and Z. Ran, *Subvarieties of generic hypersurfaces in any variety*, Math. Proc. Cambr. Phil. Soc., **130** (2001), no. 2, 259–268.
- [CR] H. Clemens, Z. Ran, *On the geometric genus of subvarieties of generic hypersurfaces*, preprint math.AG/0204256.
- [E1] L. Ein, *Subvarieties of generic complete intersections*, Invent. Math. **94** (1988), 163–169.
- [E2] L. Ein, *Subvarieties of generic complete intersections II*, Math. Ann. **289** (1991), 465–471.
- [L] S. Lang, *Hyperbolic and Diophantine analysis*, Bull. Amer. Math. Soc. (N.S.) **14** (1986), no. 2, 159–205.
- [P] G. Pacienza, *Rational curves on general projective hypersurfaces*, preprint math.AG/0010037, to appear in the J. of Alg. Geometry.
- [V1] C. Voisin, *On a conjecture of Clemens on rational curves on hypersurfaces*, J. of Diff. Geometry **44** (1996), 200–214.
- [V2] C. Voisin, *A correction on "A conjecture of Clemens on rational curves on hypersurfaces"*, J. of Diff. Geometry **49** (1998), 601–611.
- [X1] G. Xu, *Subvarieties of general hypersurfaces in projective space*, J. of Differential Geom. **39** (1994), no. 1, 139–172.
- [X2] G. Xu, *Divisors on generic complete intersection in projective space*, Trans. A.M.S. **348** (1996), 2725–2736.

Gianluca PACIENZA

Institut de Mathématiques de Jussieu

Université Pierre et Marie Curie

4, Place Jussieu, F-75252 Paris CEDEX 05 - FRANCE

e-mail: [paciENZA@math.jussieu.fr](mailto:paciENZA@math.jussieu.fr)

*current address:*

Department of Mathematics

Ohio State University

100 Mathematics Building

231 West 18th Avenue

Columbus, OH 43210-1174 - U.S.A.

e-mail: [paciENZA@math.ohio-state.edu](mailto:paciENZA@math.ohio-state.edu)